Samuel and the second second



THIE ON THE DEATHER OF AN IERRIOR

TO ALL TO WHOM THESE PRESENTS SHALL COME;

Hioneer Hi-Bred International, Inc.

Increase, there has been presented to the

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED, PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE RIGHT TO EXCLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR MPORTING IT, OR EXPORTING IT, OR CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE QVE PURPOSES, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT VARIETY THEREFROM, TO THE EXTENT DED BY THE PLANT VARIETY PROTECTION ACT. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

CORN

'PHKW3'

In Testimony Mycrost, I have hereunto set my hand and caused the seal of the Hunt Hariety Hrotection Office to be affixed at the City of Washington, D.C. this twenty-ninth day of November in the year of our Lord one thousand nine hundred and ninety-six.

REPRODUCE LOCALLY. Include form number and date on all	I reproductions		FORM APPROVED - OMB NO. 0581-00		
U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE SCIENCE DIVISION - PLANT VARIETY PROTECTION OF	The following statements are made in accordance with the Privacy Ac 1974 (5 U.S.C. 552a).				
APPLICATION FOR PLANT VARIETY PROTECTION	Application is required in order to determine if a plant variety protec certificate is to be issued 17 U.S.C. 2421). Information is held confiden				
(Instructions and information collection burden statem	ent on reverse	until certificate is issued (7 U.	S.C. 2426).		
NAME OF APPLICANT(S) (as it is to appear on the Certificate)		TEMPORARY DESIGNATION OR EXPERIMENTAL NUMBER	3. VARIETY NAME		
Pioneer Hi-Bred International	. Inc.	EX ENIMENTAL NOMBER	РНК W 3		
4. ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP Code, and Coun	ntry)	6. TELEPHONE (include area code)	FOR OFFICIAL USE ONLY		
Research & Product Development 7301 NW 62nd Avenue, PO Box 8	t	515/270-3300	9500209		
Johnston, Iowa 50131-0085		6. FAX (include area code) 515/253-2125	May 16, 1995		
7. GENUS AND SPECIES NAME	8. FAMILY NAM	E (Botanical)	FIUNG AND EXAMINATION FEE:		
Zea mays	Gra	mineae	[1:2325 ⁹⁹ +125		
9. CROP KIND NAME (Common name)	.1		DATE		
Corn			5/16/95/06/01/		
10. IF THE APPLICANT NAMED IS NOT A "PERSON", GIVE FORM OF ORGANIZA	TION (corporation,)	partnership, association, etc.) (Common name)	CERTIFICATION FEE:		
Corporation			J: 300.00		
11. IF INCORPORATED, GIVE STATE OF INCORPORATION		12. DATE OF INCORPORATION	E DATE		
Iowa -		May 6, 1926	14/13/96		
Pioneer Hi-Bred International Research & Product Development 7301 NW 62nd Avenue, PO Box 85 Johnston, IA 50131-0085 16. CHECK APPROPRIATE BOX FOR EACH ATTACHMENT SUBMITTED (Follow in	t 5	sel	515/270-3300 16. FAX (include area code) 515/253-2125		
a. C Exhibit A. Origin and Breeding History of the Variety b. 反 Exhibit B. Statement of Distinctness c. 反 Exhibit C. Objective Description of the Variety d. 反 Exhibit D. Additional Description of the Variety					
e. 🔀 Exhibit E. Statement of the Basis of the Applicant's Ownership					
 Youcher Sample (2,600 viable untreated seeds or, for tuber propagated Filing and Examination Fee (\$2,450), made payable to "Treasurer of the 			tained in a public repository)		
17. DOES THE APPLICANT SPECIFY THAT SEED OF THIS VARIETY BE SOLD BY V	VARIETY NAME ON		ection 83(a) of the Plant Variety Protection Act)?		
18. DOES THE APPLICANT SPECIFY THAT SEED OF THIS VARIETY BE LIMITED A	AS TO NUMBER OF	19. IF "YES" TO ITEM 18, WHICH CLASS	SES OF PRODUCTION BEYOND BREEDER SEED?		
GENERATIONS? YES NO		FOUNDATION REGIST	FOUNDATION REGISTERED CERTIFIED		
20. HAS THE VARIETY OR A HYBRID PRODUCED FROM THE VARIETY BEEN RELI	EASED, USED, OFF	ERED FOR SALE, OR MARKETED IN THE U.S. C	OR OTHER COUNTRIES?		
	and the second second				
21. The applicant(s) declare that a viable sample of basic seed of the variety will be applicable, or for a tuber propagated variety a tissue culture will be deposited in the undersigned applicant(s) is(are) the owner(s) of this sexually reproduced or the content of the sexually reproduced or the content of the content of the sexually reproduced or the content of the content	n a public repository tuber propagated pla	and maintained for the duration of the certificat ant variety, and believe(s) that the variety is new	e.		
Section 41, and is entitled to protection under the provisions of Section 42 of the					
Applicant(s) is(are) informed that false representation herein can jeopardize prote SIGNATURE OF APPLICANT (Owner(s))		penalties. GNATURE OF APPLICANT (Owner(s))			
		Rue O. M. Sar	etus		
NAME (Please print or type)	, i	AME (Please print or type)			
Pioneer Hi-Bred International,	Inc.	Dr. Bruce D. McBra	tnev		
CAPACITY OR TITLE DATE		APACITY OR TITLE Herbicide Research M	Resistant		
20.470					

14A. Exhibit A. Origin and Breeding History

Pedigree: PHN82/PHN63)DXA023116

Pioneer Line PHKW3, Zea mays L., a yellow corn inbred, was developed by Pioneer Hi-Bred International, Inc. from the single cross PHN82 X PHN63 using the pedigree method of breeding. The progenitors of PHKW3 are proprietary inbred lines of Pioneer Hi-Bred International, Inc. Selfing and selection were practiced within the above F1 cross for 2 generations in the development of PHKW3 at North Platte, NE. During line development, crosses were made to inbred testers for the purpose of estimating the line's combining ability. Yield trials were grown at North Platte, NE as well as other Pioneer research stations. After initial testing, additional hybrid combinations have been evaluated and subsequent generations of the line have been grown and hand-pollinated with observations made for uniformity.

PHKW3 has shown uniformity and stability for all traits as described in Exhibit C - "Objective Description of Variety". It has been self-pollinated and ear-rowed 6 generations with careful attention paid to uniformity of plant type to assure genetic homozygousity and phenotypic stability. The line has been increased both by hand and in isolated fields with continued observations for uniformity.

No variant traits have been observed or are expected in PHKW3.

The criteria used in the selection of PHKW3 were yield, both per se and in hybrid combinations; kernel size, especially important in production; ability to germinate in adverse conditions; number of tillers, especially important in production because having numerous tillers increases hybrid production costs spent on detasseling; disease and insect resistance; pollen yield; tassel size; pollen shed duration.

The inbred PHN63 traces to the public lines SRS303, I205, IDT, MINN49, WF9, OH43, OH40B, A48, and M13.

³M3 1/3/96

DEVELOPMENTAL HISTORY FOR PHKW3

Season	/Year	Inbreeding	Level
Summer	1985	F0	
Winter	1986	F1	
Summer	1986	F2*	
Summer	1988	F3	
Winter	1989	F4	
Summer	1989	F5	
Summer	1990	F6	
Summer	1991	F7	
Summer	1992	F8*	*

^{*}PHKW3 was selfed and selected through F2 generation.

**PHKW3 was selfed and ear-rowed from F3 through F8 generations.

Exhibit B. Distinctness Statement

PHKW3 is most similar to the Pioneer Hi-Bred International, Inc. proprietary inbred line PHR31 (PVP Certificate No. 9200090). The kernel rows of PHKW3 are indistinct whereas those of PHR31 are distinct. PHKW3 also differs from PHR31 in pollen shed rating (8.0 versus 5.5 rating), tassel branch angle (55.0 versus 73.0 degrees), leaf angle (37.5 versus 28.2 degrees), percent of medium round kernels (47.0 versus 25.2 percent), and cob color (white versus red) (see Attachments A and B).

PHKW3 has higher yield, higher moisture at grain harvest and higher test weight than PHR31. PHKW3 is slightly more susceptible to root and stalk lodging than PHR31. PHKW3 has better brittle stalk resistance compare to PHR31.

ATTACHMENT A - PHKW3 PIONEER CORN DESCRIPTION SYSTEM SUMMARY REPORT

Year/Trait	Mean	Standard Deviation	Environments	Replicates
93 Pollen Score	n/a	n/a	0	0
94 Pollen Score	n/a	n/a	0	0
95 Pollen Score	8.00	1.41	2	2
Pollen Score	8.00	1.41	2	2
93 Tassel Branch Angle	45.00	0.00	2	2
94 Tassel Branch Angle	45.00	0.00	2	2
95 Tassel Branch Angle	75.00	21.21	2	2
Tassel Branch Angle	55.00	18.17	6	6
93 Leaf Angle	40.00	0.00	2	2
94 Leaf Angle	35.00	7.07	2	2
95 Leaf Angle	37.50	3.54	2	2
Leaf Angle	37.50	4.18	6	6
93 Kernel % Medium Rounds	n/a	n/a	0	0
94 Kernel % Medium Rounds	48.50	2.12	2	2
95 Kernel % Medium Rounds	45.50	3.54	2	2
Kernel % Medium Rounds	47.00	2.94	4	4
93 Cob Color	White		2	2
94 Cob Color	White		2	2 .
95 Cob Color	White		2	2
Cob Color	White		6	6

ATTACHMENT B - PHR31 PIONEER CORN DESCRIPTION SYSTEM SUMMARY REPORT

Year/Trait	Mean	Standard Deviation	Environments	Replicates
93 Pollen Score	n/a	n/a	0	0
94 Pollen Score	n/a	n/a	0	0
95 Pollen Score	5.50	3.54	2	2
Pollen Score	5.50	3.54	2	2
93 Tassel Branch Angle	77.50	3.54	2	2
94 Tassel Branch Angle	45.00	0.00	1	1
95 Tassel Branch Angle	82.50	10.61	2	2
Tassel Branch Angle	73.00	16.81	5	5
93 Leaf Angle	27.50	3.54	2	2
94 Leaf Angle	25.00	0.00	1	1
95 Leaf Angle	30.50	6.36	2	2
Leaf Angle	28.20	4.32	5	5
93 Kernel % Medium Rounds	n/a	n/a	0	0
94 Kernel % Medium Rounds	27.00	0.00	2	2
95 Kernel % Medium Rounds	23.50	7.78	2	2
Kernel % Medium Rounds	25.25	4.92	4	4
93 Cob Color	Red		2	2
94 Cob Color	Red		2	2
95 Cob Color	Red		2	2
Cob Color	Red		6	6

EXHIBIT C (Corn; Maize)

United States Department of Agriculture, Agricultural Marketing Service Science Division, Plant Variety Protection Office National Agricultural Library Building, Room 500 Beltsville, MD 20705

OBJECTIVE DESCRIPTION OF VARIETY CORN (Zea mays L.)

Name of Appli		International,	, Inc.	Variety Seed Source	Variety Name or Te	mporary Designation
		.D. No., City, State, Zi	•		FOR OFFICIAL USE	
		venue, PO Box 131-0085 USA	85		PVPO Number	9500209
whole numbers Traits designa	Place the appropriate number that describes the varietal characters typical of this inbred variety in the spaces below. Right justify whole numbers by adding leading zeroes if necessary. Completeness should be striven for to establish an adequate variety description. Traits designated by a '*' are considered necessary for an adequate variety description and must be completed.					
01=Light 02=Medium 03=Dark (04=Very [Green n Green	06=Pale Yellow 07=Yellow 08=Yellow-Orange	11=Pink 12=Light Red 13=Cherry Red 14=Red	18=Colorless 19=White	21=Buff 22=Tan 23=Brown 24=Bronze	(Describe)
STANDARD INBRED CHOICES (Use the most similar (in background and maturity) of these to make comparisons based on grow-out trial data): Yellow Dent Families: Family Members Co109, ND246, C13, Iowa5125, P39, 2132 B14 CM105, A632, B64, B68 Oh7, T232 B37 B37, B76, H84 W117, W153R Popcorn: B73 N192, A679, B73, NC268 W182BN SG1533, 4722, HP301, HP7211 C103 Mo17, Va102, Va35, A682						
Oh43 WF9	A619, MS71, H9 W64A, A554, A6	99, Va26	White Dent CI66, H1	: 05, Ky228	Pipecorn: Mo15W, Mo16	W. Mo24W

COMMENTS

Color choice noted as a 26 indicates this trait was observed and recorded as green.

Data for Items 1, 3, 4, 5, 6, 7a, 7b, 8, and 9 is based primarily on a maximum of 4 reps from Johnston, Iowa, grown in 1993 and 1994, plus description information from the maintaining station.

1 mype- (accessed in the second in Comment of			Standar	d Inbred Name	
1. TYPE: (describe intermediate types in Comments section):	1 7 7		<u>B73</u>		
* <u>2</u> 1=Sweet 2=Dent 3=Flint 4=Flour 5=Pop 6=Ornamenta	il 7=Pipecoi	rn			
2. REGION WHERE DEVELOPED IN THE U.S.A.:			Standar	d Seed Source	
* <u>2</u> 1=Northwest 2=Northcentral 3=Northeast 4=Southeast	5=Southcer	ntral	AD93058	077	
6=Southwest 7=Other					
			!	•	
3. MATURITY (In Region of Best Adaptability; show Heat Unit form	ula in "Commen	ts" section	ļ		
DAYS HEAT UNITS			DAYS	HEAT UNITS	
* 67 1410.0 From emergence to 50% of plants in silk	Ξ.		6	<u>9 1557.0</u>	
* 66 1380.0 From emergence to 50% of plants in poll	.en		6	<u>9 1552.0</u>	
6131.0 From 10% to 90% pollen shed			-	5 119.0	
* From 50% silk to optimum edible quality	7				
From 50% silk to harvest at 25% moistur	re			-	
					
4 27 227	~. <u> </u>				
4. PLANT:	Standard	Sample		Standard Sample	е
	Deviation			Deviation Size	
* 224.0 cm Plant Height (to tassel tip)	<u>12.37</u>	<u>200</u>	238.0	<u>15.62</u> <u>150</u>	
* 83.0 cm Ear Height (to base of top ear node)	9.29	200	99.0	<u>8.33</u> <u>150</u>	
14.0 cm Length of Top Ear Internode	0.50	20	16.0	<u>1.73</u> <u>15</u>	
0.0 Average Number of Tillers	0.50	200	0.0	<u>0.00</u> 150	
1.0 Average Number of Ears per Stalk	0.00	200	1.0	0.00 150	
2 Anthocyanin of Brace Roots: 1=Absent 2=Faint	3=Moderate	4=Dark	4		
	•		_		
5 . T. T. T.	G	G1	_	G33 G3	
5. LEAF:	Standard	-		Standard Sample	е
	Deviation			Deviation Size	
* 9.0 cm Width of Ear Node Leaf	0.96		9.0	<u>0.58</u> <u>15</u>	
* <u>78.0</u> cm Length of Ear Node Leaf	2.52	_20	82.0	2.52 15	
* <u>5</u> Number of leaves above top ear	0.00	20	<u>6</u>	<u>0.58</u> <u>15</u>	
38 Degrees Leaf Angle (measure from 2nd leaf above	5.00	20	<u>12</u>	<u>2.89</u> <u>15</u>	
ear at anthesis to stalk above leaf					
02 Leaf Color (Munsell code 5GY 4/6)			<u>03</u> (Muns	ell Code <u>5GY 3/4</u>)	
$\underline{1}$ Leaf Sheath Pubescence (Rate on scale from 1=none to	9=like peach	fuzz)	<u>1</u>		
5 Marginal Waves (Rate on scale from 1=none to 9=many)			<u>4</u>		
6 Longitudinal Creases (Rate on scale from 1=none to 9	=many)		<u>8</u>	•	
	= 5				
5 m 60m	GE	G 1	_	Gh	_
6. TASSEL:		Sample		Standard Sample	е
16.17	Deviation	Size		Deviation Size	
* 43 Number of Primary Lateral Branches	16.17	_ 20 ^	9	<u>1.00</u> <u>15</u>	
45.0 Branch Angle from Central Spike	0.00	_20	8	<u>2.89</u> <u>15</u>	
* <u>28.0</u> cm Tassel Length (from top leaf collar to tassel	_2.36	_20	25.0	<u>1.15</u> <u>15</u>	
tip)			7		
χ Pollen Shed (rate on scale from 0=male sterile to 9=he	eavy shed)		<u>.</u> .		
07 Anther Color (Munsell code 10 Y 8 /8)			<u>07</u> (Muns	ell code <u>10YR 8/6</u>)
26 Glume Color (Munsell code 5GY 4/8) green			<u>26</u> (Muns	ell code <u>5GY 4/8</u>)	90
· · · · · · · · · · · · · · · · · · ·			<u>1</u>		•
$\underline{1}$ Bar Glumes (Glume Bands): 1=Absent 2=Present			_		
$\underline{1}$ Bar Glumes (Glume Bands): 1 =Absent 2 =Present			_		

Application Variety Data Page 2	2		Standar	d Inbred Data	
7a. EAR (Unhusked Data): Ol OSI Silk Color (3 days after emergence) (Munsell code O1 Fresh Husk Color (25 days after 50% silking) (21 Dry Husk Color (65 days after 50% silking) (Mu	Munsell code 5GY		01 (Mun	/0 sell code 2.5 sell code 5GY sell code 10Y	6/6)
1 Position of Ear at Dry Husk Stage: 1=Upright 5 Husk Tightness (Rate on scale from 1=very loose to 3 Husk extension (at harvest): 1=Short (ears exp 3=Long (8=10 cm beyond ear tip) 4=Very Long	2=Horizontal 3=I 9=very tight) posed) 2=Medium	Pendent	1 7 3	seii code <u>101</u>	
7b. EAR (Husked Ear Data):	Standard Deviation	Sample Size		Standard Deviation	Sample Size
* 13.0 cm Ear Length	0.96	20	14.0	0.58	<u>15</u>
* 46.0 mm Ear Diameter at mid-point	1.41	20	45.0	3.46	<u>15</u>
127.0 gm Ear Weight	19.02	_20	132.0	41.62	<u>15</u>
20 Number of Kernel Rows	1.00	_20	<u>17</u>	1.00	<u>15</u>
$\underline{1}$ Kernel Rows: 1=Indistinct 2=Distinct			2		
$\underline{2}$ Row Alignment: 1=Straight 2=Slightly Curved 3	=Spiral		<u>1</u>		
* <u>15.0</u> cm Shank Length	4.03	20	9.0	1.15	<u>15</u>
3. KERNEL (Dried):	Standard	Sample		Standard	Sample
	Deviation	Size		Deviation	Size
12.0 mm Kernel Length	0.50		11.0	1.15	_15
7.0 mm Kernel Width	0.50	20	8.0	0.58	_15
5.0 mm Kernel Thickness	0.50	_20	5.0	1.15	<u>15</u>
49.0 % Round Kernels (Shape Grade)	2.12	_10_	25.0	<u>17.68</u>	_15
1 Aleurone Color Pattern: 1=Homozygous 2=Segregovers (*) 07 Aleurone Color (Munsell code 2.5Y 7/12) * 07 Hard Endosperm color (Munsell code 2.5Y 8/12) * 3 Endosperm Type: 1=Sweet (Sul) 2=Extra Sweet (4=High Amylose Starch 5=Waxy 7=High Lysine 8=Super Sweet (Sul)	sh2) 3=Normal S Starch 6=High P		1	nsell code <u>2.</u>	
10=Other	3.21	<u>20</u>	26.0	4.04	_15
COB:	Standard Deviation	Sample Size		Standard Deviation	Sample Size
27.0 mm Cob Diameter at mid-point	_0.50	20	27.0	1.53	_15
19 Cob Color (Munsell code 2.5Y 9/2)		_		nsell code 10	

Application Variety Data

Page 2

Standard Inbred Data

10. DISSASS RNSISTANCE (Sate from 1 (sect susceptible) to 9 (most resistant); leave blank if not tested; leave Race or Strain Options blank if polygenic); A. Leaf Blights, Wilts, and Local Infection Diseases Anthracnose Leaf Blight (Colletotrichum graminicols) 6 Common Smut (Puccinia sorghi) Common Smut (Usrilago maydis) 5 Fyrespot (Kabatiolia zone) 4 Gray Leaf Spot (Genzopra zesa-maydis) Helminthosporium Leaf Spot (Bipolaris zelecia) Northern Leaf Blight (Esserchilum turcicum) Race 8 Southern Leaf Blight (Sisolaris gaydis) Stewart's Wilt (Eyminia stewartii) Other (Specify) 9. Systemic Diseases Corn Lethal Necrosis (MCNV and MDNV) Head Smut (Sphacelotheca roiling) Maize Chlorotic Mottle Wirus (MNV) Maize Chlorotic Mottle Wirus (MNV) Maize Chlorotic Mottle Wirus (MNV) Sorghum Downy Mildew of Corn (Peronosclerospora sorghi) Other (Specify) C. Stalk Rots Anthracnose Stalk Rot (Colletotrichum graminicols) Diplodia Stalk Rot (Stemocarpella maydis) Pusarium Stalk Rot (Stemocarpella meydis) Pusarium Stalk Rot (Stemocarpella meydis) Pusarium Stalk Rot (Stemocarpella meydis) Pusarium Sar and Kernel Rot (Ampergillus flavus) Oliplodia Sar Rot (Stemocarpella maydis) Pusarium Sar and Kernel Rot (Ampergillus flavus) Oliplodia Sar Rot (Stemocarpella maydis) Pusarium Sar and Kernel Rot (Chlorotic Desar) Glibberella Ear Rot (Glibberella zese) Other (Specify) Application Varievy Dam	Application Variety Data	Page 3	Standard Inbred Data
if not tested; leave Race or Strain Options blank if polygonic): A. Leaf Blights, Wilts, and Local Infection Diseases Anthracnose Leaf Blight (Colletotrichum graminicols) 5 Common Rust (Puccinia sorghi) 6 Common Rust (Statiago maydis) 5 Evespot (Eabtiells asses) 6 Goss's Wilt (Clavibacter michiganense spp. mebraskense) 7 Gray Leaf Spot (Corcospora reac-maydis) 8 Helminthosporium Leaf Spot (Bipolaria seicols) 8 Northern Leaf Blight (Spolaria spydis) 8 Nouthern Leaf Blight (Spolaria spydis) 8 Nace 9 Northern Leaf Blight (Spolaria spydis) 8 Southorn Rust (Puccinia polysors) 8 Stewart's Wilt (Erwinia stewartii) 9 Other (Specify) 8 Systemic Diseases Corn Lethal Necrosis (MCMV and MEMV) 9 Head Smut (Sphacelothaca reiliana) 9 Maize Chicrotic Dearf Virus (MDV) 9 Maize Chicrotic Mottle Virus (MDV) 9 Maize Chicrotic Mottle Virus (MDV) 9 Maize Chicrotic Mottle Virus (MDV) 9 Sorghum Downy Wildew of Corn (Peronosclerospora sorghi) 9 Other (Specify) C. Stalk Rots Anthracnose Stalk Rot (Colletotrichum graminicols) 9 Diplodia Stalk Rot (Stemocarpella maydis) 1 Pusarium Stalk Rot (Unsarium moniliforme) 1 Gibberella Stalk Rot (Stemocarpella maydis) 1 Diplodia Ear Rot (Stemocarpella maydis) 2 Pusarium Ear and Kernel Rot (Maspergillus flavus) 1 Ojplodia Ear Rot (Stemocarpella maydis) 2 Fissarium Ear and Kernel Rot (Maspergillus flavus) 1 Ojplodia Ear Rot (Gibberella zeae) 1 Other (Specify) 2 Other (Specify) 3 Other (Specify) 5 Other (Specify) 1 Other (Specify) 1 Other (Specify)	10 DISEASE RESISTANCE (Rate from 1 (most suscen	ntible) to 9 (most registant). leave blank	
Anthracnose Leaf Blight (Colletotrichum graminicola) 6 Common Rust (Puccinia sorghi) Common Smut (Ustilago maydis) 6 Eyespot (Kabatiella zeae) 6 Goses's Wilt (Claythacter michiganense spp. nebraskense) 7 Cray Leaf Spot (Corcespora zeae-maydis) Helminthosporium Leaf Spot (Bipolaris zeicola) Race 1 Race 1 Race 2 Race Southern Leaf Blight (Experchilum Lurcicum) Race 2 Southern Rust (Puccinia polyaora) Stewart's Wilt (Erwinia stewartii) Other (Specify) B. Systemic Diseases Corn Lethal Necrosis (MCMV and MUMV) Head Smut (Sphacelotheca reiliana) Maize Chlorotic Mottle Virus (MCMV) Sorghum Downy Mildew of Corn (Peronosclerospora sorghi) Other (Specify) C. Stalk Rote Anthracnose Stalk Rot (Colletotrichum graminicola) Diplodis Stalk Rot (Stenccarpella maydis) Fusarium Stalk Rot (Gibberella zeae) Other (Specify) D. Ear and Kernel Rote Aspergillus Ear and Kernel Rot (Aspergillus flavus) Diplodia Ear Rot (Stenccarpella maydis) Fusarium Ear and Kernel Rot (Pusarium moniliforme) Gibberella Ear Rot (Gibberella zeae) Other (Specify) 6 Other (Specify) 6 Other (Specify)			
6 Common Rust (Puccinia sorghi) Common Smut (Ustliago naydis) 6 Exyspot (Rabaticila zeas) 4 Goss's Wilt (Clavibacter michiganomse spp. nebraskense) 7 Gray Leaf Spot (Cercospora zease-maydis) 8 Helminthosporium Leaf Spot (Epipolaria gezicola) 8 Northern Leaf Blight (Exeserbhium turcicum) 9 Northern Leaf Blight (Exeserbhium turcicum) 9 Southern Leaf Blight (Exeserbhium turcicum) 9 Southern Rust (Puccinia polysora) 9 Stewart's Wilt (Erwinia stewartii) 9 Other (Specify) 9 Diplodia Stalk Not (Stenocarpella maydis) 9 Systemic Diseases 9 Corn Lethal Necrosis (MCMV and MDMV) 9 Head Smut (Sphacelotheca reilians) 9 Maize Chlorotic Dwarf Virus (MCMV) 9 Maize Chlorotic Mostle Virus (MCMV) 9 Maize Chlorotic Mostle Virus (MCMV) 9 Sorghun Downy Mildew of Corn (Peronosclerospora sorghi) 9 Other (Specify) 9 C. Stalk Rots 9 Anthracmose Stalk Rot (Colletotrichum graminicola) 9 Diplodia Stalk Rot (Stenocarpella maydis) 9 Fusarium Stalk Rot (Gibberella zeas) 9 Other (Specify) 9 D. Ear and Kernel Rot (Aspergillus flavus) 9 Diplodia Zar Rot (Stenocarpella maydis) 9 Fusarium Ear and Kernel Rot (Pusarium moniliforme) 9 Gibberella Par Rot (Gibberella zeas) 9 Other (Specify) 10 Gibberella Par Rot (Gibberella zeas) 11 Other (Specify) 12 Other (Specify) 13 Other (Specify) 14 Other (Specify) 15 Other (Specify) 16 Other (Specify) 17 Other (Specify) 18 Other (Specify) 19 Other (Specify) 19 Other (Specify) 10 Other (Specify) 10 Other (Specify) 11 Other (Specify) 12 Other (Specify) 13 Other (Specify) 14 Other (Specify) 15 Other (Specify) 16 Other (Specify) 17 Other (Specify) 18 Other (Specify) 19 Other (Specify) 19 Other (Specify) 10 Other (Specify) 10 Other (Specify) 10 Other (Specify) 11 Other (Specify) 12 Other (Specify) 13 Other (Specify) 14 Other (Specify) 15 Other (Specify) 16 Other (Specify) 17 Other (Specify) 18 Other (Specify) 19 Other (Specify) 19 Other (Specify) 10 Other (Specify) 10 Other (Specify) 10 Other (Specify) 11 Other (Specify) 12 Other (Specify) 13 Other (Specify) 14 Other (Specify) 15 Other (Specify) 16 Other (Specify) 16	A. Leaf Blights, Wilts, and Local Infection	Diseases	
Common Smut (Ustilago maydis) 6 Eyespot (Rabaticila zeae) 4 Goas's Wilt (Calvibater michiganense spp. nebraskense) 7 Gray Leaf Spot (Cercospora zeae-maydis) 8 Helminthosporium Leaf Spot (Ejeolaria zeicola) 9 Northern Leaf Blight (Exserohium turcicum) 9 Race 9 Southern Leaf Blight (Saserohium turcicum) 9 Race 9 Southern Rust (Puccinia polysora) 9 Stewart's Wilt (Erwinia stewartii) 9 Other (Specify) 9 Diplodia Stalk Rot (Colletotrichum graminicola) 9 Maize Chiorotic Mottle Virus (MCMV) 9 Maize Chiorotic Mottle Virus (MCMV) 9 Maize Chiorotic Mottle Virus (MCMV) 9 Sorghum Downy Mildew of Corn (Peronosclerospora sorghi) 9 Other (Specify) 9 Other (Specify) 9 C. Stalk Rots 9 Anthracnose Stalk Rot (Colletotrichum graminicola) 9 Diplodia Stalk Rot (Gibberella zeae) 9 Other (Specify) 9 D. Bar and Kernel Rots 9 Aspergillus Ear and Kernel Rot (Aspergillus flavus) 9 Diplodia Par Rot (Stenocarpella maydis) 9 Fusarium Ear and Kernel Rot (Pusarium moniliforme) 9 Gibberella Ear Rot (Stenocarpella maydis) 9 Fusarium Ear and Kernel Rot (Pusarium moniliforme) 9 Gibberella Ear Rot (Gibborella zeae) 9 Other (Specify) 10 Other (Specify) 11 Other (Specify) 12 Other (Specify) 13 Other (Specify) 15 Other (Specify) 16 Gibborella Ear Rot (Gibborella zeae) 17 Other (Specify) 17 Other (Specify) 18 Other (Specify) 19 Other (Specify) 19 Other (Specify) 19 Other (Specify) 10 Other (Specify) 10 Other (Specify) 10 Other (Specify) 11 Other (Specify) 12 Other (Specify) 13 Other (Specify) 15 Other (Specify) 16 Other (Specify) 17 Other (Specify) 18 Other (Specify) 19 Other (Specify) 19 Other (Specify) 10 Other (Specify) 10 Other (Specify) 10 Other (Specify) 11 Other (Specify) 12 Other (Specify) 13 Other (Specify) 14 Other (Specify) 15 Other (Specify) 15 Other (Specify) 16 Other (Specify) 17 Other (Specify) 18 Other (Specify) 19 Other (Specify) 19 Other (Specify) 19 Other (Specify) 19 Other (Specify) 10 Other (Specify) 10 Other (Specify) 11 Other (Specify) 12 Other (Specify) 13 Other (Specify) 14 Other (Specify) 15 Other (Specify) 15 Other (Specify)	_ Anthracnose Leaf Blight (<u>Colletotrichum</u>	graminicola)	_
Common Smut (Ustilago maydis) 6 Eyespot (Rabeticila zeae) 4 Goss's Wilt (Clavibacter michiganemse spp. nebraskense) 7 Gray Leaf Spot (Cercospora zeae-maydis) 8 Helminthosporium Leaf Spot (Bipolaria zeicola) 8 Helminthosporium Leaf Spot (Bipolaria zeicola) 8 Northern Leaf Blight (Exserohium turcicum) 8 Aace 9 Southern Leaf Blight (Bipolaria maydis) 8 Southern Reaf (Puccinia polysoxa) 9 Stewart's Wilt (Exvinia stewartii) 9 Other (Specify) 9 Diplodia Stalk Rot (Stenocarpella maydis) 8 Systemic Diseases Corn Lethal Necrosis (MCMV and MDMV) 9 Head Smut (Sphacelotheca reiliana) 9 Maize Chlorotic Dwarf Virus (MCMV) 9 Maize Chlorotic Mottle Virus (MCMV) 9 Maize Chlorotic Mottle Virus (MCMV) 9 Maize Chard Mosaic Virus (MCMV) 9 Sorghum Downy Mildew of Corn (Peronosclerospora sorghi) 9 Other (Specify) C. Stalk Rots Anthracnose Stalk Rot (Colletotrichum graminicola) 1 Diplodia Stalk Rot (Stenocarpella maydis) 1 Fusarium Stalk Rot (Fusarium moniliforme) 1 Gibberella Stalk Rot (Gibberella zeae) 1 Other (Specify) 1 Diplodia Bar Rot (Stenocarpella maydis) 1 Pusarium Ear and Kernel Rot (Pusarium moniliforme) 1 Gibberella Ear Rot (Stenocarpella maydis) 1 Pusarium Ear and Kernel Rot (Pusarium moniliforme) 1 Gibberella Ear Rot (Stenocarpella maydis) 1 Pusarium Ear and Kernel Rot (Pusarium moniliforme) 1 Gibberella Ear Rot (Stenocarpella maydis) 1 Pusarium Ear and Kernel Rot (Pusarium moniliforme) 1 Gibberella Ear Rot (Stenocarpella maydis) 2 Pusarium Ear and Kernel Rot (Pusarium moniliforme) 3 Gibberella Ear Rot (Gibberella zeae) 4 Other (Specify) 1 Other (Specify) 1 Other (Specify) 1 Other (Specify)	6 Common Rust (<u>Puccinia sorghi</u>)		<u>6</u>
4 Goss's Wilt (Clavibacter michiganense spp. nebraskense) 4 Gray Leaf Spot (Cercospora reae maydis) 5 Northern Leaf Spot (Epolaris reicola) Race 5 Northern Leaf Blight (Exserchilum turcicum) Race 6 Southern Rast (Puccinia polysora) 7 Southern Rust (Puccinia polysora) 8 Stewart's Wilt (Erwinia stewartii) 9 Other (Specify) 8 Systemic Diseases 9 Corn Lethal Necrosis (MCMV and MDMV) 9 Head Smut (Sphacelotheca reiliana) 9 Maize Chlorotic Dwarf Virus (MDV) 9 Maize Chlorotic Dwarf Virus (MDV) 9 Maize Dwarf Mosaic Virus (MDW) 9 Sorghum Downy Mildew of Corn (Peronosclerospora sorghi) 9 Other (Specify) 9 C. Stalk Rots 9 Anthracnose Stalk Rot (Colletotrichum graminicola) 9 Diplodia Stalk Rot (Stenocarpella maydis) 9 Fusarium Stalk Rot (Gibberella geae) 9 Other (Specify) 9 D. Ear and Kernel Rots 9 Aspergillus Ear and Kernel Rot (Aspergillus flavus) 9 Diplodia Ear Rot (Stenocarpella maydis) 9 Fusarium Ear and Kernel Rot (Pusarium moniliforme) 9 Gibberella Ear Rot (Gibberella geae) 9 Other (Specify) 1 Other (Specify) 1 Other (Specify) 2 Diplodia Ear Rot (Gibberella geae) 9 Other (Specify) 1 Other (Specify) 2 Other (Specify) 2 Other (Specify) 3 Other (Specify) 4 Other (Specify) 5 Other (Specify) 6 Other (Specify) 6 Other (Specify) 7 Other (Specify) 7 Other (Specify) 8 Other (Specify) 9 Other (Specify)	_ Common Smut (<u>Ustilago maydis</u>)		<u> </u>
Goss's Wilt (Clavibacter michiganense spp. nebraskense) 2 3 4 4 5 5 5 5 5 5 5 5	6 Eyespot (<u>Kabatiella zeae</u>)		<u>4</u>
Gray Leaf Spot (Cercospora reae-maydis)	4 Goss's Wilt (Clavibacter michiganense sp	p. nebraskense)	
Southern Leaf Blight (Sxserohilum turcicum) Race 2 Race 3 Race 4 Southern Rust (Puccinia polysora) Southern Rust (Puccinia polysora) 3 Stewart's Wilt (Erwinia stewartii) 4 Other (Specify) 4 Other (Specify) 5 Systemic Diseases 5 Systemic Diseases 6 Corn Lethal Necrosis (MCMV and MDMV) 5 Systemic Diseases 7 Systemic Diseases 8 Systemic Diseases 8 Systemic Diseases 8 Systemic Diseases 9	4 Gray Leaf Spot (Cercospora zeae-maydis)		
4 Southern Leaf Blight (Bipolaris maydis) Race	Helminthosporium Leaf Spot (Bipolaris ze	eicola) Race	Race
Southern Rust (Puccinia polysora) Stewart's Wilt (Erwinia stewartii) Other (Specify) B. Systemic Diseases Corn Lethal Necrosis (MCMV and MDMV) Head Smut (Sphacelotheca reiliana) Maize Chlorotic Dwarf Virus (MDMV) Maize Chlorotic Mostle Virus (MCMV) Sorghum Downy Mildew of Corn (Peronosclerospora sorghi) Other (Specify) C. Stalk Rots Anthracnose Stalk Rot (Colletotrichum graminicola) Diplodia Stalk Rot (Stenocarpella maydis) Fusarium Stalk Rot (Gibberella zeae) Other (Specify) D. Ear and Kernel Rots Aspergillus Ear and Kernel Rot (Aspergillus flavus) Diplodia Ear Rot (Stenocarpella maydis) Fusarium Ear and Kernel Rot (Pusarium moniliforme) Gibberella Ear Rot (Gibberella zeae) Other (Specify) Gibberella Ear Rot (Gibberella zeae)	5 Northern Leaf Blight (Exserohilum turcio	rum) Race	<u>3</u> Race
Stewart's Wilt (Erwinia stewartii) Other (Specify) B. Systemic Diseases Corn Lethal Necrosis (NCMV and MDMV) Head Smut (Sphacelotheca reiliana) Maize Chlorotic Dwarf Virus (MDW) Maize Chlorotic Mottle Virus (MCMV) Maize Chlorotic Mottle Virus (MDMV) Strain A Sorghum Downy Mildew of Corn (Peronosclerospora sorghi) Other (Specify) C. Stalk Rots Anthracnose Stalk Rot (Colletotrichum graminicola) Diplodia Stalk Rot (Stenocarpella maydis) Fusarium Stalk Rot (Gibberella zeae) Other (Specify) D. Ear and Kernel Rots Aspergillus Ear and Kernel Rot (Aspergillus flavus) Diplodia Ear Rot (Stenocarpella maydis) Fusarium Ear and Kernel Rot (Pusarium moniliforme) Gibberella Ear Rot (Stenocarpella maydis) Fusarium Ear and Kernel Rot (Pusarium moniliforme) Gibberella Ear Rot (Gibberella zeae) Other (Specify) Other (Specify) Gibberella Ear Rot (Gibberella zeae) Other (Specify)	4 Southern Leaf Blight (Bipolaris maydis)	Race	2 Race
Displodia Stalk Rot (Stenocarpella maydis) Fusarium Stalk Rot (Gibberella zeae) Other (Specify) D. Ear and Kernel Rots Aspergillus Ear Rot (Gibberella zeae) Other (Specify) D. Corn Lethal Necrosis (MCMV and MDMV) Adize Chlorotic Dwarf Virus (MDW) Maize Chlorotic Dwarf Virus (MDW) Maize Chlorotic Dwarf Virus (MDW) Sorghum Downy Mildew of Corn (Peronosclerospora sorghi) Other (Specify) C. Stalk Rots Anthracnose Stalk Rot (Colletotrichum graminicola) Diplodia Stalk Rot (Stenocarpella maydis) Fusarium Stalk Rot (Gibberella zeae) Other (Specify) D. Ear and Kernel Rots Aspergillus Ear and Kernel Rot (Aspergillus flavus) Diplodia Ear Rot (Stenocarpella maydis) Fusarium Ear and Kernel Rot (Fusarium moniliforme) Gibberella Ear Rot (Gibberella zeae) Other (Specify) D. Corn Lethal Necrosis (MCMV and MDMV) J. J	Southern Rust (<u>Puccinia</u> polysora)		_
B. Systemic Diseases Corn Lethal Necrosis (MCMV and MDMV) Head Smut (Sphacelotheca reiliana) Maize Chlorotic Dwarf Virus (MDV) Maize Chlorotic Mottle Virus (MCMV) Maize Dwarf Mosaic Virus (MCMV) Sorghum Downy Mildew of Corn (Peronosclerospora sorghi) Other (Specify) C. Stalk Rots Anthracnose Stalk Rot (Colletotrichum graminicola) Diplodia Stalk Rot (Stencarpella maydis) Fusarium Stalk Rot (Fusarium moniliforme) Gibberella Stalk Rot (Gibberella zeae) Other (Specify) D. Ear and Kernel Rots Aspergillus Ear and Kernel Rot (Aspergillus flavus) Diplodia Ear Rot (Stencarpella maydis) Fusarium Ear and Kernel Rot (Fusarium moniliforme) Gibberella Ear Rot (Stencarpella maydis) Fusarium Ear and Kernel Rot (Fusarium moniliforme) Gibberella Ear Rot (Gibberella zeae) Other (Specify) Gibberella Ear Rot (Gibberella zeae) Other (Specify)	_ Stewart's Wilt (<u>Erwinia</u> <u>stewartii</u>)		<u>4</u>
Corn Lethal Necrosis (MCMV and MDMV) Head Smut (Sphacelotheca reiliana) Maize Chlorotic Dwarf Virus (MDV) Maize Chlorotic Dwarf Virus (MDW) Maize Dwarf Mosaic Virus (MDWV) Sorghum Downy Mildew of Corn (Peronosclerospora sorghi) Other (Specify) C. Stalk Rots Anthracnose Stalk Rot (Colletotrichum graminicola) Diplodia Stalk Rot (Stenocarpella maydis) Fusarium Stalk Rot (Fusarium moniliforme) Gibberella Stalk Rot (Gibberella zeae) Other (Specify) D. Ear and Kernel Rots Aspergillus Ear and Kernel Rot (Aspergillus flavus) Diplodia Ear Rot (Stenocarpella maydis) Fusarium Ear and Kernel Rot (Fusarium moniliforme) Gibberella Ear Rot (Gibberella zeae) Other (Specify) Other (Specify) Gibberella Ear Rot (Gibberella zeae) Other (Specify) Gibberella Ear Rot (Gibberella zeae) Other (Specify) Gibberella Ear Rot (Gibberella zeae)	_ Other (Specify)		
Corn Lethal Necrosis (MCMV and MDMV) Head Smut (Sphacelotheca reiliana) Maize Chlorotic Dwarf Virus (MDV) Maize Chlorotic Mottle Virus (MCMV) Maize Dwarf Mosaic Virus (MDMV) Sorghum Downy Mildew of Corn (Peronosclerospora sorghi) Other (Specify) C. Stalk Rots Anthracnose Stalk Rot (Colletotrichum graminicola) Diplodia Stalk Rot (Stenocarpella maydis) Fusarium Stalk Rot (Fusarium moniliforme) Gibberella Stalk Rot (Gibberella zeae) Other (Specify) D. Ear and Kernel Rots Aspergillus Ear and Kernel Rot (Aspergillus flavus) Diplodia Ear Rot (Stenocarpella maydis) Fusarium Ear and Kernel Rot (Fusarium moniliforme) Gibberella Ear Rot (Gibberella zeae) Other (Specify) Other (Specify) Other (Specify) Other (Specify) Other (Specify)			
Head Smut (Sphacelotheca reiliana) Maize Chlorotic Dwarf Virus (MDV) Maize Chlorotic Mottle Virus (MDWV) Maize Dwarf Mosaic Virus (MDWV) Strain A Sorghum Downy Mildew of Corn (Peronosclerospora sorghi) Other (Specify) C. Stalk Rots Anthracnose Stalk Rot (Colletotrichum graminicola) Diplodia Stalk Rot (Stenocarpella maydis) Fusarium Stalk Rot (Fusarium moniliforme) Gibberella Stalk Rot (Gibberella zeae) Other (Specify) D. Ear and Kernel Rots Aspergillus Ear and Kernel Rot (Aspergillus flavus) Diplodia Ear Rot (Stenocarpella maydis) Fusarium Ear and Kernel Rot (Fusarium moniliforme) Gibberella Ear Rot (Gibberella zeae) Other (Specify) Other (Specify) Other (Specify)	B. Systemic Diseases		
Head Smut (Sphacelotheca reiliana) Maize Chlorotic Dwarf Virus (MDW) Maize Chlorotic Mottle Virus (MCMV) Maize Dwarf Mosaic Virus (MDMV) Sorghum Downy Mildew of Corn (Peronosclerospora sorghi) Other (Specify) C. Stalk Rots Anthracnose Stalk Rot (Colletotrichum graminicola) Diplodia Stalk Rot (Stenocarpella maydis) Fusarium Stalk Rot (Fusarium moniliforme) Gibberella Stalk Rot (Gibberella zeae) Other (Specify) D. Ear and Kernel Rots Aspergillus Ear and Kernel Rot (Aspergillus flavus) Diplodia Ear Rot (Stenocarpella maydis) Fusarium Ear and Kernel Rot (Fusarium moniliforme) Gibberella Ear Rot (Gibberella zeae) Other (Specify) Other (Specify) Other (Specify)	Corn Lethal Necrosis (MCMV and MDMV)		3
Maize Chlorotic Mottle Virus (MCMV) Maize Dwarf Mosaic Virus (MDMV) Strain A Sorghum Downy Mildew of Corn (Peronosclerospora sorghi) Other (Specify) C. Stalk Rots Anthracnose Stalk Rot (Colletotrichum graminicola) Diplodia Stalk Rot (Stenocarpella maydis) Fusarium Stalk Rot (Fusarium moniliforme) Gibberella Stalk Rot (Gibberella zeae) Other (Specify) D. Ear and Kernel Rots Aspergillus Ear and Kernel Rot (Aspergillus flavus) Diplodia Ear Rot (Stenocarpella maydis) Fusarium Ear and Kernel Rot (Fusarium moniliforme) Gibberella Ear Rot (Gibberella zeae) Other (Specify) Other (Specify) Other (Specify) Other (Specify)	Head Smut (Sphacelotheca reiliana)		_
3 Maize Dwarf Mosaic Virus (MDMV) Strain A Sorghum Downy Mildew of Corn (Peronosclerospora sorghi) Other (Specify) C. Stalk Rots Anthracnose Stalk Rot (Colletotrichum graminicola) Diplodia Stalk Rot (Stenocarpella maydis) Fusarium Stalk Rot (Fusarium moniliforme) Gibberella Stalk Rot (Gibberella zeae) Other (Specify) D. Ear and Kernel Rots Aspergillus Ear and Kernel Rot (Aspergillus flavus) Diplodia Ear Rot (Stenocarpella maydis) Fusarium Ear and Kernel Rot (Fusarium moniliforme) Gibberella Ear Rot (Gibberella zeae) Other (Specify) Other (Specify) Other (Specify)			
3 Maize Dwarf Mosaic Virus (MDMV) Strain A Sorghum Downy Mildew of Corn (Peronosclerospora sorghi) Other (Specify) C. Stalk Rots Anthracnose Stalk Rot (Colletotrichum graminicola) Diplodia Stalk Rot (Stenocarpella maydis) Fusarium Stalk Rot (Fusarium moniliforme) Gibberella Stalk Rot (Gibberella zeae) Other (Specify) D. Ear and Kernel Rots Aspergillus Ear and Kernel Rot (Aspergillus flavus) Diplodia Ear Rot (Stenocarpella maydis) Fusarium Ear and Kernel Rot (Fusarium moniliforme) Gibberella Ear Rot (Gibberella zeae) Other (Specify) Other (Specify) Other (Specify)	- Maize Chlorotic Mottle Virus (MCMV)	;	_ ,
Sorghum Downy Mildew of Corn (Peronosclerospora sorghi) Other (Specify) C. Stalk Rots Anthracnose Stalk Rot (Colletotrichum graminicola) Diplodia Stalk Rot (Stenocarpella maydis) Fusarium Stalk Rot (Fusarium moniliforme) Gibberella Stalk Rot (Gibberella zeae) Other (Specify) D. Ear and Kernel Rots Aspergillus Ear and Kernel Rot (Aspergillus flavus) Diplodia Ear Rot (Stenocarpella maydis) Fusarium Ear and Kernel Rot (Fusarium moniliforme) Gibberella Ear Rot (Gibberella zeae) Other (Specify) Other (Specify) Other (Specify)	-	Strain A	3 Strain A
C. Stalk Rots Anthracnose Stalk Rot (Colletotrichum graminicola) Diplodia Stalk Rot (Stenocarpella maydis) Fusarium Stalk Rot (Fusarium moniliforme) Gibberella Stalk Rot (Gibberella zeae) Other (Specify) D. Ear and Kernel Rots Aspergillus Ear and Kernel Rot (Aspergillus flavus) Diplodia Ear Rot (Stenocarpella maydis) Fusarium Ear and Kernel Rot (Fusarium moniliforme) Gibberella Ear Rot (Gibberella zeae) Other (Specify) Other (Specify) Other (Specify)	Sorghum Downy Mildew of Corn (Peronoscle	rospora sorghi)	
Anthracnose Stalk Rot (Colletotrichum graminicola) Diplodia Stalk Rot (Stenocarpella maydis) Fusarium Stalk Rot (Fusarium moniliforme) Gibberella Stalk Rot (Gibberella zeae) Other (Specify) D. Ear and Kernel Rots Aspergillus Ear and Kernel Rot (Aspergillus flavus) Diplodia Ear Rot (Stenocarpella maydis) Fusarium Ear and Kernel Rot (Fusarium moniliforme) Gibberella Ear Rot (Gibberella zeae) Other (Specify) Other (Specify)	-		 !
Anthracnose Stalk Rot (Colletotrichum graminicola) Diplodia Stalk Rot (Stenocarpella maydis) Fusarium Stalk Rot (Fusarium moniliforme) Gibberella Stalk Rot (Gibberella zeae) Other (Specify) D. Ear and Kernel Rots Aspergillus Ear and Kernel Rot (Aspergillus flavus) Diplodia Ear Rot (Stenocarpella maydis) Fusarium Ear and Kernel Rot (Fusarium moniliforme) Gibberella Ear Rot (Gibberella zeae) Other (Specify) Other (Specify)			· · · · · · · · · · · · · · · · · · ·
Diplodia Stalk Rot (Stenocarpella maydis) Fusarium Stalk Rot (Fusarium moniliforme) Gibberella Stalk Rot (Gibberella zeae) Other (Specify) D. Ear and Kernel Rots Aspergillus Ear and Kernel Rot (Aspergillus flavus) Diplodia Ear Rot (Stenocarpella maydis) Fusarium Ear and Kernel Rot (Fusarium moniliforme) Gibberella Ear Rot (Gibberella zeae) Other (Specify) Other (Specify)	C. Stalk Rots		
Fusarium Stalk Rot (Fusarium moniliforme) Gibberella Stalk Rot (Gibberella zeae) Other (Specify) D. Ear and Kernel Rots Aspergillus Ear and Kernel Rot (Aspergillus flavus) Diplodia Ear Rot (Stenocarpella maydis) Fusarium Ear and Kernel Rot (Fusarium moniliforme) Gibberella Ear Rot (Gibberella zeae) Other (Specify) Other (Specify)	Anthracnose Stalk Rot (Colletotrichum gr	aminicola)	
Fusarium Stalk Rot (Fusarium moniliforme) Gibberella Stalk Rot (Gibberella zeae) Other (Specify) D. Ear and Kernel Rots Aspergillus Ear and Kernel Rot (Aspergillus flavus) Diplodia Ear Rot (Stenocarpella maydis) Fusarium Ear and Kernel Rot (Fusarium moniliforme) Gibberella Ear Rot (Gibberella zeae) Other (Specify) Other (Specify)	Diplodia Stalk Rot (Stenocarpella maydis)	
D. Ear and Kernel Rots Aspergillus Ear and Kernel Rot (Aspergillus flavus) Diplodia Ear Rot (Stenocarpella maydis) Fusarium Ear and Kernel Rot (Fusarium moniliforme) Gibberella Ear Rot (Gibberella zeae) Other (Specify) ———————————————————————————————————		•	
D. Ear and Kernel Rots Aspergillus Ear and Kernel Rot (Aspergillus flavus) Diplodia Ear Rot (Stenocarpella maydis) Fusarium Ear and Kernel Rot (Fusarium moniliforme) Gibberella Ear Rot (Gibberella zeae) Other (Specify) ———————————————————————————————————	Gibberella Stalk Rot (Gibberella zeae)		
Aspergillus Ear and Kernel Rot (Aspergillus flavus) Diplodia Ear Rot (Stenocarpella maydis) Fusarium Ear and Kernel Rot (Fusarium moniliforme) Gibberella Ear Rot (Gibberella zeae) Other (Specify) ———————————————————————————————————	Other (Specify)		
Aspergillus Ear and Kernel Rot (Aspergillus flavus) Diplodia Ear Rot (Stenocarpella maydis) Fusarium Ear and Kernel Rot (Fusarium moniliforme) Gibberella Ear Rot (Gibberella zeae) Other (Specify) ———————————————————————————————————	- ·		- · · · · · · · · · · · · · · · · · · ·
Diplodia Ear Rot (Stenocarpella maydis) Fusarium Ear and Kernel Rot (Fusarium moniliforme) Gibberella Ear Rot (Gibberella zeae) Other (Specify) ———————————————————————————————————	D. Ear and Kernel Rots		
Diplodia Ear Rot (Stenocarpella maydis) Fusarium Ear and Kernel Rot (Fusarium moniliforme) Gibberella Ear Rot (Gibberella zeae) Other (Specify) ———————————————————————————————————		1 61-()	
Fusarium Ear and Kernel Rot (<u>Fusarium moniliforme</u>) Gibberella Ear Rot (<u>Gibberella zeae</u>) Other (Specify)		ius <u>riavus</u>)	_
Gibberella Ear Rot (Gibberella zeae) Other (Specify)		niliforma)	
Other (Specify)		urrroume)	
			<u> </u>
Application Variety Data Page 3 Standard Inbred Data	Other (Specify)		- .
	Application Variety Data	Page 3	Standard Inbred Data

11. INSECT RESISTANCE (Rate from 1 (most susceptible) to 9 (most resistant); (leave			
blank if not tested):			
Standard Sample		Standard	Sample
Deviation Size		Deviation	Size
_ Banks grass Mite (Oligonychus pratensis)	_		
Corn Worm (<u>Helicoverpa</u> <u>zea</u>)			
_ Leaf Feeding	_		
Silk Feeding			
mg larval wt.			
_ Ear Damage	_		
_ Corn Leaf Aphid (Rhopalosiphum maidis)			
_ Corn Sap Beetle (Carpophilus dimidiatus	_		
European Corn Borer (Ostrinia nubilalis)			
$\underline{6}$ 1 st Generation (Typically Whorl Leaf Feeding)	<u>3</u>		
4 2 nd Generation (Typically Leaf Sheath-Collar Feeding)	3		
Stalk Tunneling			
cm tunneled/plant	16.0		
Fall Armyworm (Spodoptera fruqiperda)			
Leaf Feeding			
Silk Feeding	_		
mg larval wt.	-		
Maize Weevil (Sitophilus zeamaize			
Northern Rootworm (Diabrotica barberi)	_		
Southern Rootworm (Diabrotica undecimpunctata)	- .		
Southwestern Corn Borer (Diatreaea grandiosella)	_		
Leaf Feeding			
Stalk Tunneling	_		
cm tunneled/plant			
		-	
_ Two-spotted Spider Mite (Tetranychus urticae)			
Western Rootworm (Diabrotica virgifrea virgifera)	_		
Other (Specify)	_		
		,	
12. AGRONOMIC TRAITS:			
4 Staygreen (at 65 days after anthesis) (Rate on a scale from 1=worst to	<u>3</u>		
excellent)	-		
0.0 % Dropped Ears (at 65 days after anthesis)	0.0		
% Pre-anthesis Brittle Snapping			
% Pre-anthesis Root Lodging			
Post-anthesis Root Lodging (at 65 days after anthesis)			
5650.0 Kg/ha Yield of Inbred Per Se (at 12-13% grain moisture)	4680.0		
13. MOLECULAR MARKERS: (0=data unavailable; 1=data available but not supplied;	2=data supp	olied)	
$\underline{1}$ Isozymes $\underline{0}$ RFLP's $\underline{0}$ RAPD's			
			• .
COMMENTS (eg. state how heat units were calculated, standard inbred seed source, and/or whaere data was collect	ed. Continue in I	Exhibit D);:	

ADDITIONAL DESCRIPTION OF PHKW3. INBRED PER SE YIELD TEST COMPARISON OF PHKW3 AND PHR31 EVALUATED OVER THREE YEARS. EXHIBIT D.

SIG	DRP EAR ABS	100.0 99.6 2 4 0.4			100.0 99.6 2 4 0.4
11 96	BAR PLT ABS	96.3 95.2 12 12 1.1 650	989 96.8 6.6 6.8	95.9 97.3 4 1.4	94.9 95.9 26 1.0 .601
SIG #	GRN APP ABS	4.9 4.5 7.0 1.0 889			4.0 4.7 4.0 1.0 8.8 9.8
= 5%	BRT STK ABS	99.4 99.3 7 0.0 779.	100.0 96.0 8 8 4.0 * 067*	96.7 98.6 3 1.9 689	99.2 97.7 18 20 1.4
+	STK LDG ABS	93.3 97.9 12 4.5 .105			93.3 97.9 6 12 4.5
10% SIG	STA GRN ABS	4.0 3.7 11 0.3 .825	3.0	5.0 4.3 3 0.7 635	4.0 3.7 13 17 0.3
 *	RT LDG ABS	94.2 95.9 4 1.7 686		69.4 84.3 2 14.9 .346	85.9 92.1 6.1 6.1
	GDU SLK ABS	1361 1367 13 13 15 6	1381 1412 18 18 18 31	1444 1424 11 11 20 .027+	1391 1401 42 44 10 10
	GDU SHD ABS	1331 1342 13 13 15 11 13	1349 1384 18 18 35	1414 1398 12 12 16	1362 1375 43 45 13
PHKW3 PHR31	EST CNT ABS	42.8 42.0 23 33 0.8	24.0 24.9 14 0.9	20.3 23.5 13 13 3.2 .037+	31.7 32.4 50 60 0.7 .389
f #1 - f #2 -	SDG VGR ABS	4.9 4.7 17 0.2 23	5.7 4.8 12 12 0.8	2.9 4.4 8 1.5 .026+	4.7 4.7 37 43 0.0
VARIETY VARIETY	GQU /HA \$MN	104 89 10 19 15			104 89 10 19 15
	GQU /HA ABS	59.2 51.1 10 19 8.2 .166			59.2 51.1 10 19 8.2 8.2
	TST WT ABS	55.2 53.7 15 1.5			55.2 53.7 8 15 1.5
·	MST	21.4 19.7 10 19 19 1.7			21.4 19.7 10 19 1.7 1.7
	BU ACR &MN	104 89 10 19 15			104 89 10 19 15
	BU ACR ABS	94.5 81.5 10 19 13.0			94.5 81.5 10 13.0 166
	VAR #	1 2 LOCS REPS DIFF	1 2 LOCS REPS DIFF	1 2 LOCS REPS DIFF	1 2 LOCS REPS DIFF
	REGION	SUMMARY	SUMMARY	SUMMARY	4
and the	YEAR	85	£ 6	76	TOTAL SUM

CLARIFICATION OF DATA IN EXHIBITS C AND D

Please note the data presented in Exhibit C, "Objective Description of Variety," is data collected primarily at Johnston, Iowa plus description information from the maintaining station. The data in Exhibit D, "Additional Description of Variety," is data from comparisons of inbreds grown in the same tests in the adapted growing area of PHKW3.

DEFINITIONS

In the description and examples, a number of terms are used herein. In order to provide a clear and consistent understanding of the specification and claims, including the scope to be given such terms, the following definitions are provided:

BAR PLT = BARREN PLANTS. This is the percent of plants per plot that were not barren (lack ears).

BRT STK = BRITTLE STALKS. This is a measure of the stalk breakage near the time of pollination, and is an indication of whether a hybrid or inbred would snap or break near the time of flowering under severe winds. Data are presented as percentage of plants that did not snap.

BU ACR = YIELD (BUSHELS/ACRE). Actual yield of the grain at harvest adjusted to 15.5% moisture. ABS is in absolute terms and % MN is percent of the mean for the experiments in which the hybrid or inbred was grown.

DRP EAR = DROPPED EARS. This is a measure of the number of dropped ears per plot and represents the percentage of plants that did not drop ears prior to harvest.

EAR HT = EAR HEIGHT. The ear height is a measure from the ground to the top developed ear node attachment and is measured in centimeters.

EST CNT = EARLY STAND COUNT. This is a measure of the stand establishment in the spring and represents the number of plants that emerge on a per plot basis for the hybrid or inbred.

GDU SHD = GDU TO SHED. The number of growing degree units (GDUs) or heat units required for an inbred line or hybrid to have approximately 50 percent of the plants shedding pollen and is measured from the time of planting. Growing degree units are calculated by the Barger Method, where the heat units for a 24-hour period are:

The highest maximum temperature used is 86°F and the lowest minimum temperature used is 50°F. For each inbred or hybrid it takes a certain number of GDUs to reach various stages of plant development.

<u>GDU SLK = GDU TO SILK</u>. The number of growing degree units required for an inbred line or hybrid to have approximately 50 percent of the plants with silk emergence from time of planting. Growing degree units are calculated by the Barger Method as given in GDU SHD definition.

GRN APP. = GRAIN APPEARANCE. This is a 1 to 9 rating for the general quality of the shelled grain as it is harvested based on such factors as the color of the harvested grain, any mold on the grain, and any cracked grain. High scores indicate good grain quality and low scores indicate poor grain quality.

 $\underline{MST} = \underline{HARVEST MOISTURE}$. The moisture is the actual percentage moisture of the grain at harvest.

<u>PLT HT = PLANT HEIGHT</u>. This is a measure of the height of the plant from the ground to the tip of the tassel in centimeters.

RT LDG = ROOT LODGING. Root lodging is the percentage of plants that do not root lodge; plants that lean from the vertical axis at an approximately 30° angle or greater would be counted as rootlodged.

SDG VGR = SEEDLING VIGOR. This is the visual rating (1 to 9) of the amount of vegetative growth after emergence at the seedling stage (approximately five leaves). A higher score indicates better vigor and a low score indicates poorer vigor.

STA GRN = STAY GREEN. Stay green is the measure of plant health near the time of black layer formation (physiological maturity). A high score indicates better late-season plant health.

STK LDG = STALK LODGING. This is the percentage of plants that did not stalk lodge (stalk breakage) as measured by either natural lodging or pushing the stalks and determining the percentage of plants that break below the ear.

<u>TST WT = TEST WEIGHT UNADJUSTED.</u> The measure of weight of the grain in pounds for a given volume (bushel).

14E. EXHIBIT E. Statement of the Basis of Applicant's Ownership

Pioneer Hi-Bred International, Inc., Des Moines, Iowa, is the employer of the plant breeders involved in the development and evaluation of PHKW3. Pioneer Hi-Bred International, Inc. has the sole rights and ownership of PHKW3.

REPRODUCE LOCALLY. Include form number and date on all reproductions. U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE SCIENCE AND TECHNOLOGY DIVISION - PLANT VARIETY PROTECTION OFFICE	1974 (5 U.S.C. 552a) and the Pap	le in accordance with the Privacy Act of erwork Reduction Act (PRA) of 1995 .
EXHIBIT E STATEMENT OF THE BASIS OF OWNERSHIP	Application is required in order to certificate is to be issued (7 U.S.C. until certificate is issued (7 U.S.C.	o determine if a plant variety protection 2. 2421]. Information is held confidential 2426].
1. NAME OF APPLICANT(S)	2. TEMPORARY DESIGNATION OR EXPERIMENTAL NUMBER	3. VARIETY NAME
Pioneer Hi-Bred International, Inc.		РНКW3
4. ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP Code, and Country)	5. TELEPHONE (include area code)	6. FAX (include area code)
	515/270-3300	515/253-2125
7301 NW 62nd Avenue	7. PVPO NUMBER	<u> </u>
P.O. Box 0085 Johnston, IA 50131-0085	95002	209
8. Does the applicant own all rights to the variety? Mark an "X" in appropriate b	wock. It no, please explain.	XXYES NO
·		
Is the applicant (individual or company) a U.S. national or U.S. based company If no, give name of country		XXYES NO
10. Is the applicant the original breeder? If no, please answer the following:		XX YES NO
 a. If original rights to variety were owned by individual(s): ls (are) the original breeder(s) a U.S. national(s)? If no, give name of c 	ountry	
		TYES NO
 b. If original rights to variety were owned by a company: ls the original breeder(s) U.S. based company? If no, give name of course 	untry	
11. Additional explantion on ownership (If needed, use reverse for extra space):		
PLEASE NOTE: .		
Plant variety protection can be afforded only to owners (not licensees) who meet o	ne of the following criteria:	
If the rights to the variety are owned by the original breeder, that person must of a country which affords similar protection to nationals of the U.S. for the sar		UPOV member country, or national
If the rights to the variety are owned by the company which employed the originationals of a UPOV member country, or owned by nationals of a country which genus and species.		
3. If the applicant is an owner who is not the original breeder, both the original breeder.	eder and the applicant must mee	t one of the above criteria.
The original breeder may be the individual or company who directed final breed definition.	ing. See Section 41(a)(2) of th	e Plant Variety Protection Act for

Public reporting burden for this collection of information is estimated to average 10 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Department of Agriculture, Clearance Officer, OIRM, AG Box 7630, Jamie L. Whitten Building, Washington, D.C. 20250. When replying, refer to OMB No. 0581-0055 and form number in your letter.

Under the PRA of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

The U.S. Department of Agriculture (USDA) prohibits discrimination in its programs on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, and marital or familial status. (Not all prohibited bases apply to all programs). Persons with disabilities who require alternative means for communication of program information (braille, large print, audiotape, etc.) should contact the USDA Office of Communications at (202) 720-2791.

To file a complaint, write the Secretary of Agriculture, U.S. Department of Agriculture, Washington, D.C. 20250, or call (202) 720-7327 (voice) or (202) 720-1127 (TDD). USDA is an equal employment opportunity employer.